

Artwork No.			Colours Used
Customer	Somex Pharma	■	Black
Description	Rivaroxaban FC Tablets_10 mg_Package Leaflet		
Market	UK		
Language	English		
Size	144 x 544 mm		
Min. Font Size			
Version No.	07		
Pharma code			
Date	07.01.2026		
Software	Adobe Illustrator cs6		

144 mm

**Package leaflet: Information for the user**  
**Rivaroxaban 10 mg film-coated tablets**  
**Rivaroxaban**

**Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

**What is in this leaflet**

1. What Rivaroxaban tablet is and what it is used for
2. What you need to know before you take Rivaroxaban tablet
3. How to take Rivaroxaban tablet
4. Possible side effects
5. How to store Rivaroxaban tablet
6. Contents of the pack and other information

**1. What Rivaroxaban tablet is and what it is used for**

Rivaroxaban tablet contains the active substance rivaroxaban and is used in adults to

- prevent blood clots in the veins after a hip or knee replacement operation. Your doctor has prescribed this medicine for you because after an operation you are at an increased risk of getting blood clots.
- treat blood clots in the veins of your legs (deep vein thrombosis) and in the blood vessels of your lungs (pulmonary embolism), and to prevent blood clots from re-occurring in the blood vessels of your legs and/or lungs.

Rivaroxaban tablet belongs to a group of medicines called antithrombotic agents. It works by blocking a blood clotting factor (factor Xa) and thus reducing the tendency of the blood to form clots.

**2. What you need to know before you take Rivaroxaban tablet**

**Do not take Rivaroxaban tablet**

- if you are allergic to rivaroxaban or any of the other ingredients of this medicine (listed in section 6)
- if you are bleeding excessively
- if you have a disease or condition in an organ of the body that increases the risk of serious bleeding (e.g. stomach ulcer, injury or bleeding in the brain, recent surgery of the brain or eyes)
- if you are taking medicines to prevent blood clotting (e.g. warfarin, dabigatran, apixaban or heparin), except when changing anticoagulant treatment or while getting heparin through a venous or arterial line to keep it open
- if you have a liver disease which leads to an increased risk of bleeding
- if you are pregnant or breast-feeding

**Do not take Rivaroxaban tablet and tell your doctor if any of these apply to you.**

**Warnings and precautions**

Talk to your doctor or pharmacist before taking Rivaroxaban tablet.

**Take special care with Rivaroxaban tablet**

- if you have an increased risk of bleeding, as could be the case in situations such as:
  - moderate or severe kidney disease, since your kidney function may affect the amount of medicine that works in your body
  - if you are taking other medicines to prevent blood clotting (e.g. warfarin, dabigatran, apixaban or heparin), when changing anticoagulant treatment or while getting heparin through a venous or arterial line to keep it open (see section "Other medicines and Rivaroxaban tablet")
  - bleeding disorders
  - very high blood pressure, not controlled by medical treatment
  - diseases of your stomach or bowel that might result in bleeding, e.g. inflammation of the bowels or stomach, or inflammation of the oesophagus (gullet), e.g. due to gastroesophageal reflux disease (disease where stomach acid goes upwards into the oesophagus)
  - a problem with the blood vessels in the back of your eyes (retinopathy)
  - a lung disease where your bronchi are widened and filled with pus (bronchiectasis), or previous bleeding from your lung
- if you have a prosthetic heart valve
- if you know that you have a disease called antiphospholipid syndrome (a disorder of the immune system that causes an increased risk of blood clots), tell your doctor who will decide if the treatment may need to be changed.
- if your doctor determines that your blood pressure is unstable or removal of a blood clot from your lungs is planned.

**If any of the above apply to you, tell your doctor before you take Rivaroxaban tablet.** Your doctor will decide, if you should be treated with this medicine and if you should be kept under closer observation.

**If you need to have an operation**

- it is very important to take Rivaroxaban tablet before and after the operation exactly at the times you have been told by your doctor.
- If your operation involves a catheter or injection into your spinal column (e.g. for epidural or spinal anaesthesia or pain reduction):
  - it is very important to take Rivaroxaban tablet exactly at the times you have been told by your doctor

- tell your doctor immediately if you get numbness or weakness of your legs or problems with your bowel or bladder after the end of anaesthesia, because urgent care is necessary.

**Children and adolescents**

Rivaroxaban tablet is **not recommended for people under 18 years of age**. There is not enough information on its use in children and adolescents.

**Other medicines and Rivaroxaban tablet**

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines, including medicines obtained without a prescription.

**If you are taking**

- some medicines for fungal infections (e.g. fluconazole, itraconazole, voriconazole, posaconazole), unless they are only applied to the skin
- ketoconazole tablets (used to treat Cushing's syndrome - when the body produces an excess of cortisol)
- some medicines for bacterial infections (e.g. clarithromycin, erythromycin)
- some anti-viral medicines for HIV / AIDS (e.g. ritonavir)
- other medicines to reduce blood clotting (e.g. enoxaparin, clopidogrel or vitamin K antagonists such as warfarin and acenocoumarol)
- anti-inflammatory and pain relieving medicines (e.g. naproxen or acetylsalicylic acid)
- dronedarone, a medicine to treat abnormal heart beat
- some medicines to treat depression (selective serotonin reuptake inhibitors (SSRIs) or serotonin norepinephrine reuptake inhibitors (SNRIs))

**If any of the above apply to you, tell your doctor before taking Rivaroxaban tablet,** because the effect of Rivaroxaban tablet may be increased. Your doctor will decide, if you should be treated with this medicine and if you should be kept under closer observation.

If your doctor thinks that you are at increased risk of developing stomach or bowel ulcers, he may also use a preventative ulcer treatment.

**If you are taking**

- some medicines for treatment of epilepsy (phenytoin, carbamazepine, phenobarbital)
- St John's Wort (Hypericum perforatum), a herbal product used for depression
- rifampicin, an antibiotic

**If any of the above apply to you, tell your doctor before taking Rivaroxaban tablet,** because the effect of Rivaroxaban tablet may be reduced. Your doctor will decide, if you should be treated with Rivaroxaban tablet and if you should be kept under closer observation.

**Pregnancy and breast-feeding**

Do not take Rivaroxaban tablet if you are pregnant or breast-feeding. If there is a chance that you could become pregnant, use a reliable contraceptive while you are taking Rivaroxaban tablet. If you become pregnant while you are taking this medicine, tell your doctor immediately, who will decide how you should be treated.

**Driving and using machines**

Rivaroxaban tablet may cause dizziness (common side effect) or fainting (uncommon side effect) (see section 4 "Possible side effects"). You should not drive or use machines if you are affected by these symptoms.

**Rivaroxaban tablet contains lactose and sodium**

If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicine. This medicine contains less than 1 mmol sodium (23 mg) per tablet, that is to say essentially "sodium-free".

**3. How to take Rivaroxaban tablet**

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

**How much to take**

- To prevent blood clots in the veins after a hip or knee replacement operation. The recommended dose is one tablet Rivaroxaban 10 mg once a day.
- To treat blood clots in the veins of your legs and blood clots in the blood vessels of your lungs, and for preventing blood clots from re-occurring  
After at least 6 months blood clot treatment, the recommended dose is either one 10 mg tablet once a day or one 20 mg tablet once a day. Your doctor has prescribed you Rivaroxaban tablet 10 mg once a day.

Swallow the tablet preferably with water. Rivaroxaban tablet can be taken with or without food.

If you have difficulty swallowing the tablet whole, talk to your doctor about other ways to take Rivaroxaban tablet. The tablet may be crushed and mixed with water or apple puree immediately before you take it. If necessary, your doctor may also give you the crushed Rivaroxaban tablet through a stomach tube.

**When to take Rivaroxaban tablet**

Take the tablet every day until your doctor tells you to stop. Try to take the tablet at the same time every day to help you to remember it. Your doctor will decide how long you must continue treatment.

544 mm

**To prevent blood clots in the veins after a hip or knee replacement operation:** Take the first tablet 6 - 10 hours after your operation.

If you have had a major hip operation you will usually take the tablets for 5 weeks. If you have had a major knee operation you will usually take the tablets for 2 weeks.

**If you take more Rivaroxaban tablet than you should**

Contact your doctor immediately if you have taken too many Rivaroxaban tablet. Taking too much Rivaroxaban tablet increases the risk of bleeding.

**If you forget to take Rivaroxaban tablet**

If you have missed a dose, take it as soon as you remember. Take the next tablet on the following day and then carry on taking a tablet once a day as normal.

Do not take a double dose to make up for a forgotten tablet.

**If you stop taking Rivaroxaban tablet**

Do not stop taking Rivaroxaban tablet without talking to your doctor first, because Rivaroxaban tablet prevents the development of a serious condition.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

**4. Possible side effects**

Like all medicines, Rivaroxaban tablet can cause side effects, although not everybody gets them.

Like other similar medicines (antithrombotic agents), Rivaroxaban tablet may cause bleeding which may potentially be life threatening. Excessive bleeding may lead to a sudden drop in blood pressure (shock). In some cases the bleeding may not be obvious.

**Tell your doctor immediately**, if you experience any of the following side effects:

**Signs of bleeding**

- bleeding into the brain or inside the skull (symptoms can include headache, one-sided weakness, vomiting, seizures, decreased level of consciousness, and neck stiffness. A serious medical emergency. Seek medical attention immediately!)
- long or excessive bleeding
- exceptional weakness, tiredness, paleness, dizziness, headache, unexplained swelling, breathlessness, chest pain or angina pectoris, which may be signs of bleeding.

Your doctor may decide to keep you under closer observation or change how you should be treated.

**Signs of severe skin reactions**

- spreading intense skin rash, blisters or mucosal lesions, e.g. in the mouth or eyes (Stevens- Johnson syndrome/ toxic epidermal necrolysis). The frequency of this side effect is very rare (up to 1 in 10,000).
- a drug reaction that causes rash, fever, inflammation of internal organs, blood abnormalities and systemic illness (DRESS syndrome). The frequency of this side effect is very rare (up to 1 in 10,000).

**Signs of severe allergic reactions**

- swelling of the face, lips, mouth, tongue or throat; difficulty swallowing; hives and breathing difficulties; sudden drop in blood pressure. The frequencies of these side effects are very rare (anaphylactic reactions, including anaphylactic shock; may affect up to 1 in 10,000 people) and uncommon (angioedema and allergic oedema; may affect up to 1 in 100 people).

**Overall list of possible side effects**

**Common** (may affect up to 1 in 10 people)

- reduction in red blood cells which can make the skin pale and cause weakness or breathlessness
- bleeding in the stomach or bowel, urogenital bleeding (including blood in the urine and heavy menstrual bleeding), nose bleed, bleeding in the gum
- bleeding into the eye (including bleeding from the whites of the eyes)
- bleeding into tissue or a cavity of the body (haematoma, bruising)
- coughing up blood
- bleeding from the skin or under the skin
- bleeding following an operation
- oozing of blood or fluid from surgical wound
- swelling in the limbs
- pain in the limbs
- impaired function of the kidneys (may be seen in tests performed by your doctor)
- fever
- stomach ache, indigestion, feeling or being sick, constipation, diarrhoea
- low blood pressure (symptoms may be feeling dizzy or fainting when standing up)
- decreased general strength and energy (weakness, tiredness), headache, dizziness
- rash, itchy skin
- blood tests may show an increase in some liver enzymes

**Uncommon** (may affect up to 1 in 100 people)

- bleeding into the brain or inside the skull
- bleeding into a joint causing pain and swelling
- thrombocytopenia (low number of platelets, which are cells that help blood to clot)
- allergic reactions, including allergic skin reactions
- impaired function of the liver (may be seen in tests performed by your doctor)
- blood tests may show an increase in bilirubin, some pancreatic or liver enzymes or in the number of platelets
- fainting
- feeling unwell

- faster heartbeat
- dry mouth
- hives

**Rare** (may affect up to 1 in 1,000 people)

- bleeding into a muscle
- cholestasis (decreased bile flow), hepatitis incl. hepatocellular injury (inflamed liver incl. liver injury)
- yellowing of the skin and eye (jaundice) localised swelling
- collection of blood (haematoma) in the groin as a complication of the cardiac procedure where a catheter is inserted in your leg artery (pseudoaneurysm)

**Very rare** (may affect up to 1 in 10,000 people)

- accumulation of eosinophils, a type of white granulocytic blood cells that cause inflammation in the lung (eosinophilic pneumonia)

**Not known** (frequency cannot be estimated from the available data)

- kidney failure after a severe bleeding
- bleeding in the kidney sometimes with presence of blood in urine leading to inability of the kidneys to work properly (anticoagulant-related nephropathy)
- increased pressure within muscles of the legs or arms after a bleeding, which leads to pain, swelling, altered sensation, numbness or paralysis (compartment syndrome after a bleeding)

**Reporting of side effects**

If you get any side effects, talk to your doctor or pharmacist or nurse.

This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard) or search for MHRA Yellow Card in the Google Play or Apple App Store.

By reporting side effects you can help provide more information on the safety of this medicine.

**5. How to store Rivaroxaban tablet**

Keep out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton and on each blister or bottle after EXP. The expiry date refers to the last day of that month.

This medicine does not require any special storage conditions.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

**6. Contents of the pack and other information**

**What Rivaroxaban tablet contains**

- The active substance is rivaroxaban. Each tablet contains 10 mg of rivaroxaban, respectively.
- The other ingredients are:  
Tablet core: microcrystalline cellulose (E460), croscarmellose sodium, lactose monohydrate, hypromellose 2910/5 (E464), sodium laurilsulphate, magnesium stearate. See section 2 "Rivaroxaban tablet contains lactose and sodium".  
Tablet film coat: HPMC 2910/hypromellose (E464), macrogol/PEG (E1521), titanium dioxide (E171), FD&C blue #2/indigo carmine aluminum lake (E132), FD&C blue #2/indigo carmine AL3%-5% (E132).

**What Rivaroxaban tablet looks like and contents of the pack**

Rivaroxaban 10 mg film-coated tablets are blue, round (diameter: 6 mm), biconvex and marked with a "10" on one side and plain on the other side.

They come

- in blisters in cartons of 10, 30 or 100 film-coated tablets or
- in unit dose blisters in cartons of 10 x 1 or 100 x 1 or
- in multipacks comprising 10 cartons, each containing 10 x 1 film-coated tablets.

Not all pack sizes may be marketed.

**Marketing Authorisation Holder**

Somex Pharma,  
Ilford, Essex,  
IG3 8BS, UK

**Manufacturer**

Bluepharma - Indústria Farmacêutica, S.A.  
São Martinho do Bispo  
3045-016 Coimbra, Portugal

**This leaflet was last revised in December 2025**

Artwork No.			Colours Used
Customer	Somex Pharma		Black
Description	Rivaroxaban FC Tablets_ 15 mg/20 mg_PI		
Market	UK		
Language	English		
Size	144 x 544 mm		
Min. Font Size			
Version No.	04		
Pharma code			
Date	07.01.2026		
Software	Adobe Illustrator cs6		

144 mm

Package leaflet: Information for the user

**Rivaroxaban 15 mg film-coated tablets**  
**Rivaroxaban 20 mg film-coated tablets**  
**Rivaroxaban**

**Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

**What is in this leaflet**

1. What Rivaroxaban tablet is and what it is used for
2. What you need to know before you take Rivaroxaban tablet
3. How to take Rivaroxaban tablet
4. Possible side effects
5. How to store Rivaroxaban tablet
6. Contents of the pack and other information

**1. What Rivaroxaban tablet is and what it is used for**

Rivaroxaban tablet contains the active substance rivaroxaban and is used in adults to:

- prevent blood clots in brain (stroke) and other blood vessels in your body if you have a form of irregular heart rhythm called non-valvular atrial fibrillation.
- treat blood clots in the veins of your legs (deep vein thrombosis) and in the blood vessels of your lungs (pulmonary embolism), and to prevent blood clots from re-occurring in the blood vessels of your legs and/or lungs.
- Rivaroxaban tablet is used in children and adolescents below 18 years and with a body weight of 30 kg or more to:
  - treat blood clots and prevent re- occurrence of blood clots in the veins or in the blood vessels of the lungs, following initial treatment of at least 5 days with injectable medicines used to treat blood clots.

Rivaroxaban tablet belongs to a group of medicines called antithrombotic agents. It works by blocking a blood clotting factor (factor Xa) and thus reducing the tendency of the blood to form clots.

**2. What you need to know before you take Rivaroxaban tablet**

**Do not take Rivaroxaban tablet**

- if you are allergic to rivaroxaban or any of the other ingredients of this medicine (listed in section 6)
- if you are bleeding excessively
- if you have a disease or condition in an organ of the body that increases the risk of serious bleeding (e.g. stomach ulcer, injury or bleeding in the brain, recent surgery of the brain or eyes)
- if you are taking medicines to prevent blood clotting (e.g. warfarin, dabigatran, apixaban or heparin), except when changing anticoagulant treatment or while getting heparin through a venous or arterial line to keep it open.
- if you have a liver disease which leads to an increased risk of bleeding
- if you are pregnant or breast-feeding

**Do not take Rivaroxaban tablet and tell your doctor** if any of these apply to you.

**Warnings and precautions**

Talk to your doctor or pharmacist before taking Rivaroxaban Tablet.

**Take special care with Rivaroxaban Tablet**

- if you have an increased risk of bleeding, as could be the case in situations such as:
  - severe kidney disease, since your kidney function may affect the amount of medicine that works in your body
  - if you are taking other medicines to prevent blood clotting (e.g. warfarin, dabigatran, apixaban or heparin), when changing anticoagulant treatment or while getting heparin through a venous or arterial line to keep it open (see section "Other medicines and Rivaroxaban Tablet")
  - bleeding disorders
  - very high blood pressure, not controlled by medical treatment
  - diseases of your stomach or bowel that might result in bleeding, e.g. inflammation of the bowels or stomach, or inflammation of the oesophagus (gullet), e.g. due to gastroesophageal reflux disease (disease where stomach acid goes upwards into the oesophagus)
  - a problem with the blood vessels in the back of your eyes (retinopathy)
  - a lung disease where your bronchi are widened and filled with pus (bronchiectasis), or previous bleeding from your lung
- if you have a prosthetic heart valve
- if you know that you have a disease called antiphospholipid syndrome (a disorder of the immune system that causes an increased risk of blood clots), tell your doctor who will decide if the treatment may need to be changed.
- if your doctor determines that your blood pressure is unstable or another treatment or surgical procedure to remove the blood clot from your lungs is planned

**If any of the above apply to you, tell your doctor** before you take Rivaroxaban Tablet. Your doctor will decide, if you should be treated with this medicine and if you should be kept under closer observation.

**If you need to have an operation**

- it is very important to take Rivaroxaban Tablet before and after the operation exactly at the times you have been told by your doctor.
- If your operation involves a catheter or injection into your spinal column (e.g. for epidural or spinal anaesthesia or pain reduction):
  - it is very important to take Rivaroxaban Tablet before and after the injection or removal of the catheter exactly at the times you have been told by your doctor
  - tell your doctor immediately if you get numbness or weakness of your legs or problems with your bowel or bladder after the end of anaesthesia, because urgent care is necessary.

**Children and adolescents**

Rivaroxaban Tablet is **not recommended for children with a body weight below 30 kg**. There is not enough information on its use in children and adolescents.

**Other medicines and Rivaroxaban Tablet**

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines, including medicines obtained without a prescription.

**If you are taking**

- some medicines for fungal infections (e.g. fluconazole, itraconazole, voriconazole, posaconazole), unless they are only applied to the skin
- ketoconazole tablets (used to treat Cushing's syndrome - when the body produces an excess of cortisol)

- some medicines for bacterial infections (e.g. clarithromycin, erythromycin)
- some anti-viral medicines for HIV / AIDS (e.g. ritonavir)
- other medicines to reduce blood clotting (e.g. enoxaparin, clopidogrel or vitamin K antagonists such as warfarin and acenocoumarol)
- anti-inflammatory and pain relieving medicines (e.g. naproxen or acetylsalicylic acid)
- dronedarone, a medicine to treat abnormal heart beat
- some medicines to treat depression (selective serotonin reuptake inhibitors (SSRIs) or serotonin norepinephrine reuptake inhibitors (SNRIs))

**If any of the above apply to you, tell your doctor** before taking Rivaroxaban Tablet, because the effect of Rivaroxaban Tablet may be increased. Your doctor will decide, if you should be treated with this medicine and if you should be kept under closer observation.

If your doctor thinks that you are at increased risk of developing stomach or bowel ulcers, he may also use a preventative ulcer treatment.

**If you are taking**

- some medicines for treatment of epilepsy (phenytoin, carbamazepine, phenobarbital)
- St John's Wort (Hypericum perforatum), a herbal product used for depression
- rifampicin, an antibiotic

**If any of the above apply to you, tell your doctor** before taking Rivaroxaban Tablet, because the effect of Rivaroxaban Tablet may be reduced. Your doctor will decide, if you should be treated with Rivaroxaban Tablet and if you should be kept under closer observation.

**Pregnancy and breast-feeding**

Do not take Rivaroxaban Tablet if you are pregnant or breast-feeding. If there is a chance that you could become pregnant, use a reliable contraceptive while you are taking Rivaroxaban Tablet. If you become pregnant while you are taking this medicine, tell your doctor immediately, who will decide how you should be treated.

**Driving and using machines**

Rivaroxaban Tablet may cause dizziness (common side effect) or fainting (uncommon side effect) (see section 4, "Possible side effects"). You should not drive or use machines if you are affected by these symptoms.

**Rivaroxaban Tablet contains lactose and sodium**

If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicine.

This medicine contains less than 1 mmol sodium (23 mg) per tablet that is to say essentially "sodium- free".

**3. How to take Rivaroxaban Tablet**

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

You must take Rivaroxaban Tablet together with a meal. Swallow the tablet(s) preferably with water.

If you have difficulty swallowing the tablet whole, talk to your doctor about other ways to take Rivaroxaban Tablet. The tablet may be crushed and mixed with water or apple puree immediately before you take it. This mixture should be immediately followed by food.

If necessary, your doctor may also give you the crushed Rivaroxaban Tablet through a stomach tube.

**How much to take**

**Adults**

- To prevent blood clots in (stroke) and other blood vessels in your body the recommended dose is one tablet Rivaroxaban Tablet 20 mg once a day.
  - If you have kidney problems, the dose may be reduced to one tablet Rivaroxaban Tablet 15 mg once a day.
  - If you need a procedure to treat blocked blood vessels in your heart (called a percutaneous coronary intervention - PCI with an insertion of a stent), there is limited evidence to reduce the dose to one tablet Rivaroxaban Tablet 15 mg once a day (or to one tablet Rivaroxaban Tablet 10 mg once a day in case your kidneys are not working properly) in addition to an antiplatelet medicinal product such as clopidogrel.
- To treat blood clots in the veins of your legs and blood clots in the blood vessels of your lungs, and for preventing blood clots from re-occurring
  - The recommended dose is one tablet Rivaroxaban Tablet 15 mg twice a day for the first 3 weeks. For treatment after 3 weeks, the recommended dose is one tablet Rivaroxaban Tablet 20 mg once a day.
  - After at least 6 months blood clot treatment your doctor may decide to continue treatment with either one 10 mg tablet once a day or one 20 mg tablet once a day.
  - If you have kidney problems and take one tablet Rivaroxaban Tablet 20 mg once a day, your doctor may decide to reduce the dose for the treatment after 3 weeks to one tablet Rivaroxaban Tablet 15 mg once a day if the risk for bleeding is greater than the risk for having another blood clot.

**Children and adolescents**

The dose of Rivaroxaban tablet depends on the body weight, and will be calculated by the doctor.

- The recommended dose for children and adolescents with a body weight between 30 kg and less than 50 kg is one Rivaroxaban 15mg tablet once a day.
- The recommended dose for children and adolescents with a body weight of 50 kg or more is one Rivaroxaban 20 mg tablet once a day. Take each Rivaroxaban tablet dose with a drink (e.g. water or juice) during a meal. Take the tablets every day at approximately the same time. Consider setting an alarm to remind you. For parents or caregivers: please observe the child to ensure the full dose is taken.

As the Rivaroxaban tablet dose is based on body weight it is important to keep scheduled doctor's visits because the dose may need to be adjusted as the weight changes. **Never adjust the dose of Rivaroxaban tablet by yourself**. The doctor will adjust the dose if necessary.

Do not split the tablet in an attempt to provide a fraction of a tablet dose. If a lower dose is required please use the alternative presentation of Rivaroxaban.

For children and adolescents who are unable to swallow tablets whole, please use alternative presentation of Rivaroxaban.

**If you spit up the dose or vomit**

- less than 30 minutes after you have taken Rivaroxaban tablet, take a new dose.
- more than 30 minutes after you have taken Rivaroxaban tablet, do not take a new dose. In this case, take the next Rivaroxaban tablet dose at the usual time. Contact the doctor if you repeatedly spit up the dose or vomit after taking Rivaroxaban tablet.

**When to take Rivaroxaban Tablet**

Take the tablet(s) every day until your doctor tells you to stop.

Try to take the tablet(s) at the same time every day to help you to remember it. Your doctor will decide how long you must continue treatment.

To prevent blood clots in the brain (stroke) and other blood vessels in your body:

If your heart beat needs to be restored to normal by a procedure called cardioversion, take Rivaroxaban Tablet at the times your doctor tells you.

**If you take more Rivaroxaban Tablet than you should**

**Contact your doctor immediately** if you have taken too many Rivaroxaban Tablets. Taking too much Rivaroxaban Tablet increases the risk of bleeding.

**If you forget to take Rivaroxaban Tablet****Adults, children and adolescents:**

- If you are taking one 20 mg tablet or one 15 mg tablet once a day and have missed a dose, take it as soon as you remember. Do not take more than one tablet in a single day to make up for a forgotten dose. Take the next tablet on the following day and then carry on taking one tablet once a day.

**Adults:**

- If you are taking one 15 mg tablet twice a day and have missed a dose, take it as soon as you remember. Do not take more than two 15 mg tablets in a single day. If you forget to take a dose you can take two 15 mg tablets at the same time to get a total of two tablets (30 mg) on one day. On the following day you should carry on taking one 15 mg tablet twice a day.

**If you stop taking Rivaroxaban Tablet**

Do not stop taking Rivaroxaban Tablet without talking to your doctor first, because Rivaroxaban Tablet treats and prevents serious conditions.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

**4. Possible side effects**

Like all medicines, Rivaroxaban Tablet can cause side effects, although not everybody gets them.

Like other similar medicines (antithrombotic agents), Rivaroxaban Tablet may cause bleeding which may potentially be life threatening. Excessive bleeding may lead to a sudden drop in blood pressure (shock). In some cases the bleeding may not be obvious.

**Tell your doctor immediately** if you or the child experience any of the following side effects:

**Signs of bleeding**

- Bleeding into the brain or inside the skull (symptoms can include headache, one-sided weakness, vomiting, seizures, decreased level of consciousness, and neck stiffness. A serious medical emergency. Seek medical attention immediately!)
- long or excessive bleeding
- exceptional weakness, tiredness, paleness, dizziness, headache, unexplained swelling, breathlessness, chest pain or angina pectoris, which may be signs of bleeding.

Your doctor may decide to keep you under closer observation or change how you should be treated.

**Signs of severe skin reactions**

- spreading intense skin rash, blisters or mucosal lesions, e.g. in the mouth or eyes (Stevens- Johnson syndrome/toxic epidermal necrolysis). The frequency of this side effect is very rare (up to 1 in 10,000).
- a drug reaction that causes rash, fever, inflammation of internal organs, blood abnormalities and systemic illness (DRESS syndrome). The frequency of this side effect is very rare (up to 1 in 10,000).

**Signs of severe allergic reactions**

- swelling of the face, lips, mouth, tongue or throat; difficulty swallowing; hives and breathing difficulties; sudden drop in blood pressure. The frequencies of these side effects are very rare (anaphylactic reactions, including anaphylactic shock; may affect up to 1 in 10,000 people) and uncommon (angioedema and allergic oedema; may affect up to 1 in 100 people).

**Overall list of possible side effects****Common** (may affect up to 1 in 10 people)

- reduction in red blood cells which can make the skin pale and cause weakness or breathlessness
- bleeding in the stomach or bowel, urogenital bleeding (including blood in the urine and heavy menstrual bleeding), nose bleed, bleeding in the gum
- bleeding into the eye (including bleeding from the whites of the eyes)
- bleeding into tissue or a cavity of the body (haematoma, bruising)
- coughing up blood
- bleeding from the skin or under the skin
- bleeding following an operation
- oozing of blood or fluid from surgical wound
- swelling in the limbs
- pain in the limbs
- impaired function of the kidneys (may be seen in tests performed by your doctor)
- fever
- stomach ache, indigestion, feeling or being sick, constipation, diarrhoea
- low blood pressure (symptoms may be feeling dizzy or fainting when standing up)
- decreased general strength and energy (weakness, tiredness), headache, dizziness
- rash, itchy skin
- blood tests may show an increase in some liver enzymes

**Uncommon** (may affect up to 1 in 100 people)

- bleeding into the brain or inside the skull
- bleeding into a joint causing pain and swelling
- thrombocytopenia (low number of platelets, which are cells that help blood to clot)
- allergic reactions, including allergic skin reactions
- impaired function of the liver (may be seen in tests performed by your doctor)
- blood tests may show an increase in bilirubin, some pancreatic or liver enzymes or in the number of platelets
- fainting
- feeling unwell
- faster heartbeat
- dry mouth
- hives

**Rare** (may affect up to 1 in 1,000 people)

- bleeding into a muscle
- cholestasis (decreased bile flow), hepatitis incl. hepatocellular injury (inflamed liver incl. liver injury)
- yellowing of the skin and eye (jaundice)
- localised swelling
- collection of blood (haematoma) in the groin as a complication of the cardiac procedure where a catheter is inserted in your leg artery (pseudoaneurysm)

**Very rare** (may affect up to 1 in 10,000 people)

- accumulation of eosinophils, a type of white granulocytic blood cells that cause inflammation in the lung (eosinophilic pneumonia)

**Not known** (frequency cannot be estimated from the available data)

- kidney failure after a severe bleeding
- Bleeding in kidney sometimes with presence of blood in urine leading to inability of the kidneys to work properly (anticoagulant-related nephropathy)
- increased pressure within muscles of the legs or arms after a bleeding, which leads to pain, swelling, altered sensation, numbness or paralysis (compartment syndrome after a bleeding)

**Side effects in children and adolescents**

In general, the side effects observed in children and adolescents treated with Rivaroxaban tablets were similar in type to those observed in adults and were primarily mild to moderate in severity.

Side effects that were observed more often in children and adolescents:

**Very common** (may affect more than 1 in 10 people)

- headache
- fever
- nose bleeding
- vomiting

**Common** (may affect up to 1 in 10 people)

- raised heartbeat
- blood tests may show an increase in bilirubin (bile pigment)
- thrombocytopenia (low number of platelets which are cells that help blood to clot)
- heavy menstrual bleeding

**Uncommon** (may affect up to 1 in 100 people)

- blood tests may show an increase in a subcategory of bilirubin (direct bilirubin, bile pigment)

**Reporting of side effects**

If you get any side effects, talk to your doctor or pharmacist or nurse.

This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at:

[www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard) or search for MHRA Yellow Card in the Google Play or Apple App Store.

By reporting side effects you can help provide more information on the safety of this medicine.

**5. How to store Rivaroxaban Tablet**

Keep out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton and on each blister or bottle after EXP.

The expiry date refers to the last day of that month.

This medicine does not require any special storage conditions.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

**6. Contents of the pack and other information****What Rivaroxaban Tablet contains**

- The active substance is rivaroxaban. Each tablet contains 15 mg or 20 mg of rivaroxaban, respectively.
- The other ingredients are:  
Tablet core: microcrystalline cellulose (E460), croscarmellose sodium, lactose monohydrate, hypromellose 2910/5 (E464), sodium laurilsulphate, magnesium stearate. See section 2 "Rivaroxaban Tablet contains lactose and sodium".  
Tablet film coat 15 mg: HPMC 2910/hypromellose (E 464), titanium dioxide (E171) and macrogol/PEG (E1521).  
Tablet film coat 20 mg: HPMC 2910/hypromellose (E464), iron oxide red (E172), titanium dioxide (E171) and macrogol/PEG (E1521) and talc (E 553b).

**What Rivaroxaban Tablet looks like and contents of the pack**

Rivaroxaban 15 mg film-coated tablets are white, round (diameter: 6 mm), biconvex and marked with a "15" on one side and plain on the other side.

They come

- in blisters in cartons of 14, 28, 42 or 100 film-coated tablets or
- in unit dose blisters in cartons of 14 x 1 or 100 x 1 or
- in multipacks comprising 10 cartons, each containing 10 x 1 film-coated tablets.

Rivaroxaban 20 mg film-coated tablets are red, round (diameter: 6 mm), biconvex and marked with a "20" on one side and plain on the other side.

They come

- in blisters in cartons of 14, 28 or 100 film-coated tablets or
- in unit dose blisters in cartons of 14 x 1 or 100 x 1 or
- in multipacks comprising 10 cartons, each containing 10 x 1 film-coated tablets.

Not all pack sizes may be marketed.

**Marketing Authorisation Holder**

Somex Pharma,  
Ilford, Essex,  
IG3 8BS, UK

**Manufacturer**

Bluepharma - Indústria Farmacêutica, S.A.  
São Martinho do Bispo  
3045-016 Coimbra,  
Portugal

**This leaflet was last revised in December 2025**